

Worthless

SEARCH REQUEST FORM

Access DB# 60538

Scientific and Technical Information Center

Received back 3/28

Requester's Full Name: P. Pawack Examiner #: 70400 Date: 3/15/02
Art Unit: 1614 Phone Number 30 84703 Serial Number: 09/995277
Mail Box and Bldg/Room Location: 2D05 Results Format Preferred (circle): PAPER DISK E-MAIL

If more than one search is submitted, please prioritize searches in order of need.

Please provide a detailed statement of the search topic, and describe as specifically as possible the subject matter to be searched. Include the elected species or structures, keywords, synonyms, acronyms, and registry numbers, and combine with the concept or utility of the invention. Define any terms that may have a special meaning. Give examples or relevant citations, authors, etc., if known. Please attach a copy of the cover sheet, pertinent claims, and abstract.

Title of Invention: Fc Receptor Modulators


Inventors (please provide full names): Jonathan B. Baell

Earliest Priority Filing Date: 9/11/98


For Sequence Searches Only Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.

Please search methods for inhibiting Fc receptor binding of immunoglobulin

surprising administering a compound of



Ar = aryl (no heterocycles)



L = alkylene optionally with oxygen atoms

G = a bond

- CH_2

- alkylene

Thank

=> fil hcaplus
 FILE 'HCAPLUS' ENTERED AT 10:23:08 ON 21 MAR 2002
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.


FILE COVERS 1907 - 21 Mar 2002 VOL 136 ISS 12
 FILE LAST UPDATED: 19 Mar 2002 (20020319/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

CAS roles have been modified effective December 16, 2001. Please check your SDI profiles to see if they need to be revised. For information on CAS roles, enter HELP ROLES at an arrow prompt or use the CAS Roles thesaurus (/RL field) in this file.

The P indicator for Preparations was not generated for all of the CAS Registry Numbers that were added to the CAS files between 12/27/01 and 1/23/02. As of 1/23/02, the situation has been resolved. Searches and/or SDIs in the H/Z/CA/CAplus files incorporating CAS Registry Numbers with the P indicator executed between 12/27/01 and 1/23/02 may be incomplete. See the NEWS message on this topic for more information.

=>
 =>
 => d stat que 131
 L1 SCR 1840 OR 2127 OR 2050 OR 2049 OR 2048 OR 2053 OR 2052 O
 R 2051 OR 2043
 L2 STR



REP G1=(0-10) C
 REP G2=(2-5) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 GGCAT IS MCY AT 8

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 14


STEREO ATTRIBUTES: NONE

L3 0 SEA FILE=REGISTRY SSS FUL L2 NOT L1

L4 SCR 1840 OR 2127 OR 2050 OR 2049 OR 2048 OR 2053 OR 2052 O

R 2051 OR 2043

L5 STR



REP G1=(0-10) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 8

DEFAULT ECLEVEL IS LIMITED


GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 11

STEREO ATTRIBUTES: NONE

L6 STR



REP G1=(0-10) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 8

DEFAULT ECLEVEL IS LIMITED


GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 12

STEREO ATTRIBUTES: NONE


L7 STR



REP G1=(0-10) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 GGCAT IS MCY AT 8
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 12


STEREO ATTRIBUTES: NONE
 L8 STR



REP G1=(0-10) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 GGCAT IS MCY AT 8
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 14


STEREO ATTRIBUTES: NONE
 L9 STR



REP G1=(0-10) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 GGCAT IS MCY AT 8
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 14

STEREO ATTRIBUTES: NONE
 L10 STR



REP G1=(0-10) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 GGCAT IS MCY AT 8
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 14

STEREO ATTRIBUTES: NONE
 L11 (575)SEA FILE=REGISTRY SSS FUL L5 OR L6 OR L7
 L12 (2502)SEA FILE=REGISTRY SSS FUL L8 OR L9
 L13 (1035)SEA FILE=REGISTRY SSS FUL L10 NOT L4
 L14 4087 SEA FILE=REGISTRY ABB=ON PLU=ON L11 OR L12 OR L13
 L20 STR

O=C~O
6 @7 @8


N=C~N~OH
9 @10 11 12

O=C~CF3
13 @14 15

O=P~O
16 @17 @18

C~G3~G1
@19 20 21

C=C~G3~G1
@22 23 24 25



VAR G1=7/10/14/17
VAR G2=7/10/8/14/17/18/19/22

REP G3=(1-10) C

REP G5=(0-10) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 2

GGCAT IS MCY AT 4

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 28

STEREO ATTRIBUTES: NONE

L21 94 SEA FILE=REGISTRY SSS FUL L20

L22 STR

O=C~O
6 @7 @8


N=C~N~OH
9 @10 11 12

O=C~CF3
13 @14 15

O=P~O
16 @17 @18

C~G3~G1
@19 20 21

C=C~G3~G1
@22 23 24 25



VAR G1=7/10/14/17
VAR G2=7/10/8/14/17/18/19/22

REP G3=(1-10) C

REP G5=(0-10) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 2

GGCAT IS MCY AT 4

DEFAULT ECLEVEL IS LIMITED








GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 29

STEREO ATTRIBUTES: NONE

L23 STR



VAR G1=7/10/14/17

VAR G2=7/10/8/14/17/18/19/22

REP G3=(1-10) C

REP G5=(0-10) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 2

GGCAT IS MCY AT 4

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 28

STEREO ATTRIBUTES: NONE

L24 21 SEA FILE=REGISTRY SSS FUL L22 OR L23

L25 4201 SEA FILE=REGISTRY ABB=ON PLU=ON L3 OR L14 OR L21 OR L24

L26 1831 SEA FILE=HCAPLUS ABB=ON PLU=ON L25

L27 21 SEA FILE=REGISTRY ABB=ON PLU=ON FC RECEPTOR?/CN

L29 6587 SEA FILE=HCAPLUS ABB=ON PLU=ON L27 OR (FC OR F(W)C) (W) RECEPTOR?

L31 0 SEA FILE=HCAPLUS ABB=ON PLU=ON L26 AND L29

=>


=>

=> d stat que 132

L1 SCR 1840 OR 2127 OR 2050 OR 2049 OR 2048 OR 2053 OR 2052 O

R 2051 OR 2043

L2 STR



REP G1=(0-10) C

REP G2=(2-5) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 8

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 14


STEREO ATTRIBUTES: NONE

L3 0 SEA FILE=REGISTRY SSS FUL L2, NOT L1

L4, SCR 1840 OR 2127 OR 2050 OR 2049 OR 2048 OR 2053 OR 2052 O

R 2051 OR 2043

L5 STR



REP G1=(0-10) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 8

DEFAULT ECLEVEL IS LIMITED


GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 11

STEREO ATTRIBUTES: NONE

L6 STR



REP G1=(0-10) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 8

DEFAULT ECLEVEL IS LIMITED


GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 12

STEREO ATTRIBUTES: NONE


L7 STR



REP G1=(0-10) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 GGCAT IS MCY AT 8
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 12


STEREO ATTRIBUTES: NONE
 L8 STR



REP G1=(0-10) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 GGCAT IS MCY AT 8
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 14


STEREO ATTRIBUTES: NONE
 L9 STR



REP G1=(0-10) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 GGCAT IS MCY AT 8
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 14

STEREO ATTRIBUTES: NONE
 L10 STR



REP G1=(0-10) C
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 GGCAT IS MCY AT 8
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 14

STEREO ATTRIBUTES: NONE

L11 (575)SEA FILE=REGISTRY SSS FUL L5 OR L6 OR L7
 L12 (2502)SEA FILE=REGISTRY SSS FUL L8 OR L9
 L13 (1035)SEA FILE=REGISTRY SSS FUL L10 NOT L4
 L14 4087 SEA FILE=REGISTRY ABB=ON PLU=ON L11 OR L12 OR L13
 L20 STR

O=C~O
6 @7 @8


N=C~N~OH
9 @10 11 12

O=C~CF3
13 @14 15

O=P~O
16 @17 @18

C~G3~G1
@19 20 21

C=C~G3~G1
@22 23 24 25



VAR G1=7/10/14/17
VAR G2=7/10/8/14/17/18/19/22

REP G3=(1-10) C

REP G5=(0-10) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 2

GGCAT IS MCY AT 4

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 28

STEREO ATTRIBUTES: NONE

L21 94 SEA FILE=REGISTRY SSS FUL L20

L22 STR

O=C~O
6 @7 @8


N=C~N~OH
9 @10 11 12

O=C~CF3
13 @14 15

O=P~O
16 @17 @18

C~G3~G1
@19 20 21

C=C~G3~G1
@22 23 24 25



VAR G1=7/10/14/17

VAR G2=7/10/8/14/17/18/19/22

REP G3=(1-10) C

REP G5=(0-10) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 2

GGCAT IS MCY AT 4

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 29

STEREO ATTRIBUTES: NONE

L23 STR

O=C~O
6 @7 @8


N=C~N~OH
9 @10 11 12

O=C~CF3
13 @14 15

O=P~O
16 @17 @18

C~G3~G1
@19 20 21

C=C~G3~G1
@22 23 24 25



VAR G1=7/10/14/17
VAR G2=7/10/8/14/17/18/19/22

REP G3=(1-10) C

REP G5=(0-10) C

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

GGCAT IS MCY AT 2

GGCAT IS MCY AT 4

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 28

STEREO ATTRIBUTES: NONE

L24 21 SEA FILE=REGISTRY SSS FUL L22 OR L23

L25 4201 SEA FILE=REGISTRY ABB=ON PLU=ON L3 OR L14 OR L21 OR L24

L26 1831 SEA FILE=HCAPLUS ABB=ON PLU=ON L25

L28 28801 SEA FILE=REGISTRY ABB=ON PLU=ON IMMUNOGLOBULIN?

L30 151724 SEA FILE=HCAPLUS ABB=ON PLU=ON L28 OR ?IMMUNOGLOBULIN? OR IG#

L32 1 SEA FILE=HCAPLUS ABB=ON PLU=ON L26 AND L30

=>

=>

=> d ibib abs hitrn l32 1

L32 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1977:567872 HCAPLUS

DOCUMENT NUMBER: 87:167872

TITLE: Furanone derivatives as fluorogenic reagents

INVENTOR(S): Cleeland, Roy, Jr.; Grunberg, Emanuel; Leimgruber, Willy; Weigele, Manfred

PATENT ASSIGNEE(S): Hoffmann-La Roche, Inc., USA

SOURCE: U.S., 8 pp. Division of U.S. 3,969,373.

CODEN: USXXAM

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 5


PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 4045487	A	19770830	US 1976-668179	19760318
NO 7500485	A	19740906	NO 1974-485	19750214
US 3969373	A	19760713	US 1975-590655	19750626
US 4238589	A	19801209	US 1978-966527	19781204

PRIORITY APPLN. INFO.:

US 1973-338019	19730305
US 1975-590655	19750626
NO 1974-682	19740227
US 1976-668179	19760318
US 1977-801952	19770531

GI



AB Six furanones I (R = Me, Et, PhCH₂; R₁ = Ph, p-O₂NC₆H₄, p-HO₂CC₆H₄, p-HO₂CCH₂CH₂C₆H₄), useful for fluorescent labeling of biol. materials, e.g., amino acids, peptides, **immunoglobulins**, and bacteria, were prepd. from benzalacetophenones by multistep sequence. Thus, benzalacetophenone was epoxidized under basic conditions, the resultant epoxy ketone was treated with Me₃COK to cleave the epoxide ring, and the produced dione was treated with (MeO)₂CHNMe₂ to give PhCOCOCPh:CHNMe₂, which was hydrolyzed with aq. KOH, the basic soln. was acidified with HCl, and the product (I; R = H, R₁ = Ph) was methylated with MeOH to give I (R = Me, R₁ = Ph). I are themselves nonfluorescent but produce highly fluorescent substances upon reaction with primary amine-contg. compds. Six examples of fluorescent labeling, e.g., of *Hymenolepis hana*, *Escherichia coli*, and γ -globulin, with I are given.

IT 20118-38-1P 61059-47-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)
(prepn. and epoxidation of)

=>

=> d stat que 152 nos

L1 SCR 1840 OR 2127 OR 2050 OR 2049 OR 2048 OR 2053 OR 2052 O

R 2051 OR 2043

L2 STR

L3 0 SEA FILE=REGISTRY SSS FUL L2 NOT L1

L4 SCR 1840 OR 2127 OR 2050 OR 2049 OR 2048 OR 2053 OR 2052 O

R 2051 OR 2043

L5 STR

L6 STR

L7 STR

L8 STR

L9 STR

L10 STR

L11 (575) SEA FILE=REGISTRY SSS FUL L5 OR L6 OR L7

L12 (2502) SEA FILE=REGISTRY SSS FUL L8 OR L9

L13 (1035) SEA FILE=REGISTRY SSS FUL L10 NOT L4

L14 4087 SEA FILE=REGISTRY ABB=ON PLU=ON L11 OR L12 OR L13

L20 STR

L21 94 SEA FILE=REGISTRY SSS FUL L20

L22 STR

L23 STR

L24 21 SEA FILE=REGISTRY SSS FUL L22 OR L23

L25 4201 SEA FILE=REGISTRY ABB=ON PLU=ON L3 OR L14 OR L21 OR L24

L26 1831 SEA FILE=HCAPLUS ABB=ON PLU=ON L25
 L27 21 SEA FILE=REGISTRY ABB=ON PLU=ON FC RECEPTOR?/CN
 L28 28801 SEA FILE=REGISTRY ABB=ON PLU=ON IMMUNOGLOBULIN?
 L29 6587 SEA FILE=HCAPLUS ABB=ON PLU=ON L27 OR (FC OR F(W)C) (W)RECEPTO
 R?
 L30 151724 SEA FILE=HCAPLUS ABB=ON PLU=ON L28 OR ?IMMUNOGLOBULIN? OR
 IG#
 L32 1 SEA FILE=HCAPLUS ABB=ON PLU=ON L26 AND L30
 L37 31935 SEA FILE=HCAPLUS ABB=ON PLU=ON L29 OR FC OR F(W)C
 L38 1 SEA FILE=HCAPLUS ABB=ON PLU=ON L37 AND L26
 L52 1 SEA FILE=HCAPLUS ABB=ON PLU=ON L38 NOT L32

=>

=>

=> d ibib abs hitrn 152 1

L52 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1999:441770 HCAPLUS

DOCUMENT NUMBER: 131:229682

TITLE: Dye-Labeled Poly(organosiloxane) Microgels with
 Core-Shell Architecture

AUTHOR(S): Graf, Christina; Schaertl, Wolfgang; Fischer, Karl;
 Hugenberg, Norbert; Schmidt, Manfred

CORPORATE SOURCE: Institut fuer Physikalische Chemie, Universitaet
 Mainz, Mainz, 55099, Germany

SOURCE: Langmuir (1999), 15(19), 6170-6180

CODEN: LANGD5; ISSN: 0743-7463

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Poly(organosiloxane) microgels are highly cross-linked rather monodisperse
 spherical particles of radius about 10 nm. Using a functionalized silane
 comonomer, i.e., (chlorobenzyl)trimethoxysilane, model particles suitable
 for studies in colloid physics are available: photoreactive and
 fluorescent dyes can be covalently bound within the microgels to prep.
 tracers for diffusion studies using forced Rayleigh scattering (FRS) and
 fluorescence correlation spectroscopy (FCS). For the
 application as tracer particles, it is important not to influence the
 diffusion behavior by the coupled chromophores. Therefore, functionalized
 precursors with a core-shell architecture are used to minimize labeling
 effects. The photochromic dye ortho-nitrostilbene (ONS) and the
 fluorophores rhodamine B, coumarin 343, and pyrene, resp., were then
 coupled to the functionalized cores. The dye content of the labeled
 .mu.-gels strongly decreases with increasing thickness of the protective
 shell. A higher polarity of the used chromophores also lowers the dye
 content significantly, while differences in the size of the used label
 mols. are less important. The fluorescence intensity of the dye-labeled
 spheres is also influenced by the size of the protective shell which has
 been explained by differences in mobility of the labels (caging effects)
 and, at high dye concn. (thinner shell), by reabsorption.

IT 65199-97-5DP, salts or esters, reaction products with
 chlorobenzyltrimethoxysilane-methyltrimethoxysilane copolymer

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
 (dye-labeled poly(organosiloxane) microgels with core-shell
 architecture)

REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d stat que.153 nos

L1 SCR 1840 OR 2127 OR 2050 OR 2049 OR 2048 OR 2053 OR 2052 O
 R 2051 OR 2043
 L2 STR
 L3 0 SEA FILE=REGISTRY SSS FUL L2 NOT L1
 L4 SCR 1840 OR 2127 OR 2050 OR 2049 OR 2048 OR 2053 OR 2052 O
 R 2051 OR 2043
 L5 STR
 L6 STR
 L7 STR
 L8 STR
 L9 STR
 L10 STR
 L11 (575) SEA FILE=REGISTRY SSS FUL L5 OR L6 OR L7
 L12 (2502) SEA FILE=REGISTRY SSS FUL L8 OR L9
 L13 (1035) SEA FILE=REGISTRY SSS FUL L10 NOT L4
 L14 4087 SEA FILE=REGISTRY ABB=ON PLU=ON L11 OR L12 OR L13
 L20 STR
 L21 94 SEA FILE=REGISTRY SSS FUL L20
 L22 STR
 L23 STR
 L24 21 SEA FILE=REGISTRY SSS FUL L22 OR L23
 L25 4201 SEA FILE=REGISTRY ABB=ON PLU=ON L3 OR L14 OR L21 OR L24
 L26 1831 SEA FILE=HCAPLUS ABB=ON PLU=ON L25
 L27 21 SEA FILE=REGISTRY ABB=ON PLU=ON FC RECEPTOR?/CN
 L28 28801 SEA FILE=REGISTRY ABB=ON PLU=ON IMMUNOGLOBULIN?
 L29 6587 SEA FILE=HCAPLUS ABB=ON PLU=ON L27 OR (FC OR F(W)C) (W)RECEPTO
 R?
 L30 151724 SEA FILE=HCAPLUS ABB=ON PLU=ON L28 OR ?IMMUNOGLOBULIN? OR
 IG#
 L32 1 SEA FILE=HCAPLUS ABB=ON PLU=ON L26 AND L30
 L33 106 SEA FILE=HCAPLUS ABB=ON PLU=ON L26(L) (?RECEPTOR? OR ?BIND?)
 L35 152 SEA FILE=HCAPLUS ABB=ON PLU=ON L26(L) ?INHIBIT?
 L36 23 SEA FILE=HCAPLUS ABB=ON PLU=ON L33 AND L35
 L37 31935 SEA FILE=HCAPLUS ABB=ON PLU=ON L29 OR FC OR F(W)C
 L38 1 SEA FILE=HCAPLUS ABB=ON PLU=ON L37 AND L26
 L40 59 SEA FILE=HCAPLUS ABB=ON PLU=ON L26(L) (?MODULAT? OR ?CONTROL?
 OR ?REGULAT?)
 L41 5 SEA FILE=HCAPLUS ABB=ON PLU=ON L40 AND L35
 L42 26 SEA FILE=HCAPLUS ABB=ON PLU=ON L41 OR L36
 L52 1 SEA FILE=HCAPLUS ABB=ON PLU=ON L38 NOT L32
 L53 26 SEA FILE=HCAPLUS ABB=ON PLU=ON L42 NOT (L32 OR L38 OR L52)

=>

=> d ibib.abs.hitrn 153 1-26

L53 ANSWER 1 OF 26 HCAPLUS COPYRIGHT 2002 ACS
 ACCESSION NUMBER: 2001:103108 HCAPLUS
 DOCUMENT NUMBER: 134:217509
 TITLE: Inhibition of ovulation in the rat by a leukotriene B4
 receptor antagonist
 AUTHOR(S): Matousek, Markus; Mitsube, Kenrokuro; Mikuni, Masato;
 Brannstrom, Mats
 CORPORATE SOURCE: Department of Obstetrics and Gynecology, Goteborg
 University, Goteborg, S-413 45, Swed.
 SOURCE: Molecular Human Reproduction (2001), 7(1), 35-42
 CODEN: MHREFD; ISSN: 1360-9947

PUBLISHER: Oxford University Press
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The involvement of leukotriene (LT) B₄ in the ovulatory process of the rat was investigated by the use of a LTB₄-receptor antagonist (ZK 158252 = L-ANT) administered either intrabursally in vivo or to the in-vitro perfused ovary. The in-vivo expts. revealed inhibition of human chorionic gonadotropin (HCG)-induced ovulation by 500 $\mu\text{mol/L}$ L-ANT (median 5.5, 25-75% range 1.0-6.0) compared with controls (median 9.0, range 6.25-13.5). In vitro, ovulation was induced by LH (0.2 $\mu\text{g/mL}$) + 3-isobutyl-1-methylxanthine (IBMX; 0.2 mmol/L). The ovary was perfused either for 20 h, to study ovulation rate, or for 10 h to examine ovarian concns. of the ovulatory mediators matrix metalloproteinase (MMP)-2 and MMP-9, plasminogen activator (PA), prostaglandin (PG)E₂ and PGF₂.alpha.. Addn. of LH + IBMX resulted in a marked stimulation of steroid release and ovulations occurred in all ovaries (median 11.0, range 10.0-14.0). The L-ANT inhibited ovulation in a dose-dependent way (median 10.0, range 8.0-13.0 at 1 $\mu\text{mol/L}$; median 6.0, range 3.5-10.0 at 10 $\mu\text{mol/L}$; median 2.0, range 0.75-5.75 at 100 $\mu\text{mol/L}$). The intra-ovarian activity of PA was increased 1.5-fold by L-ANT (100 $\mu\text{mol/L}$), but the concns. of PGE₂ and PGF₂.alpha. remained unaltered. While no changes in MMP-9 were obsd., conversion from pro-MMP-2 to active MMP-2 was inhibited by L-ANT. These results suggest that activation of the LTB₄-receptor within the ovary is involved in the ovulatory process and that the effects of LTB₄-receptor activation are partly mediated via MMP-2.

IT 245742-21-6, ZK 158252

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(leukotriene B₄ receptor antagonist inhibition of ovulation in rats)

REFERENCE COUNT: 58 THERE ARE 58 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L53 ANSWER 2 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2000:899249 HCAPLUS

DOCUMENT NUMBER: 135:55572

TITLE: Retinoic acid (RA) receptor transcriptional activation correlates with inhibition of 12-O-tetradecanoylphorbol-13-acetate-induced ornithine decarboxylase (ODC) activity by retinoids: A potential role for trans-RA-induced ZBP-89 in ODC inhibition

AUTHOR(S): Dawson, Marcia I.; Park, Ju Rui; Chen, Guo-Quan; Chao, Wan-Ru; Dousman, Linda; Waleh, Nahid; Hobbs, Peter D.; Jong, Ling; Toll, Lawrence; Zhang, Xiao-Kun; Gu, Jian; Agadir, Anissa; Merchant, Juanita L.; Bai, Longchuan; Verma, Ajit K.; Thatcher, Scott M.; Chandraratna, Roshantha A. S.; Shroot, Braham; Hill, Donald L.

CORPORATE SOURCE: Department of Medicinal Chemistry, Molecular Medicine Research Institute, Mountain View, CA, 94043, USA

SOURCE: International Journal of Cancer (2001), 91(1), 8-21
CODEN: IJCNAW; ISSN: 0020-7136

PUBLISHER: Wiley-Liss, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Evaluation of retinoic acid receptor (RAR) subtype-selective .alpha. and .gamma. agonists and antagonists and a retinoid X receptor (RXR) class-selective agonist for efficacy at inhibiting both induction of ornithine decarboxylase (ODC) by the tumor promoter 12-O-

tetradecanoylphorbol-13-acetate (TPA) in mouse epidermis and rat tracheal epithelial cells and the appearance of papillomas in mouse epidermis treated in the 2-stage tumor initiation-promotion model indicated that (i) RXR class-selective transcriptional agonists, such as MMI 1246, were not involved in ODC inhibition; (ii) RAR-selective agonists that induce gene transcription from RA-responsive elements (RAREs) were active at low concns.; (iii) RAR-selective antagonists that bind RARs and inhibit AP-1 activation on the collagenase promoter but do not activate RAREs to induce gene transcription were less effective inhibitors; and (iv) RAR.gamma.-selective retinoid agonists were more effective inhibitors of TPA-induced ODC activity than RAR.alpha.-selective agonists. These results suggest that RARE activation has a more important role in inhibition of ODC activity than RXR activation or AP-1 inhibition and that RAR.gamma.-selective agonists would be the most useful inhibitors of epithelial cell proliferation induced by tumor promoters. The natural retinoid all-trans-RA induced expression of transcription factor ZBP-89, which represses activation of the GC box in the ODC promoter by the transcription factor Sp1.

IT 138254-19-0, MM 3986

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(retinoic acid **receptor** transcriptional activation correlates with **inhibition** of tumor promoter-induced ornithine decarboxylase activity by retinoids and role for ZBP-89 therein)

REFERENCE COUNT: 62 THERE ARE 62 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L53 ANSWER 3 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2000:307979 HCAPLUS

DOCUMENT NUMBER: 133:202727

TITLE: Identification of receptor-selective retinoids that are potent inhibitors of the growth of human head and neck squamous cell carcinoma cells

AUTHOR(S): Sun, Shi-Yong; Yue, Ping; Mao, Li; Dawson, Marcia I.; Shroot, Braham; Lamph, William W.; Heyman, Richard A.; Chandraratna, Roshantha A. S.; Shudo, Koichi; Hong, Waun K.; Lotan, Reuben

CORPORATE SOURCE: Department of Thoracic/Head and Neck Medical Oncology, The University of Texas M. D. Anderson Cancer Center, Houston, TX, 77030, USA

SOURCE: Clinical Cancer Research (2000), 6(4), 1563-1573
CODEN: CCREF4; ISSN: 1078-0432

PUBLISHER: American Association for Cancer Research

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Retinoids modulate the growth and differentiation of cancer cells presumably by activating gene transcription via the nuclear retinoic acid receptor (RAR) .alpha., .beta., and .gamma. and retinoid X receptor (RXR) .alpha., .beta., and .gamma.. We analyzed the effects of 38 RAR-selective and RXR-selective retinoids on the proliferation of 10 human head and neck squamous cell carcinoma (HNSCC) cell lines. All of these cell lines expressed constitutively all of the receptor subtypes except RAR.beta., which was detected in only two of them. Most of the RAR-selective retinoids inhibited the growth of HNSCC cells to varying degrees, whereas the RXR-selective retinoids showed very weak or no inhibitory effects. Three RAR antagonists suppressed growth inhibition by RAR-selective agonists, as well as by RAR/RXR antagonists such as 9-cis-retinoic acid. Combinations of RXR-selective and RAR-selective retinoids exhibited additive growth-inhibitory effects. Furthermore, we found that CD437, the

most potent growth-inhibitory retinoid induced apoptosis and up-regulated the expression of several apoptosis-related genes in HNSCC cells. These results indicate that: (a) retinoid receptors are involved in the growth-inhibitory effects of retinoids; (b) RXR-RAR heterodimers rather than RXR-RXR homodimer are the major mediators of growth inhibition by retinoids in HNSCC cells; and (c) induction of apoptosis can account for one mechanism by which retinoids such as CD437 inhibit the growth of HNSCC cells. Finally, these studies identified several synthetic retinoids, which are much more effective than the natural RAs and can be good candidates for chemoprevention and therapy of head and neck cancers.

IT 110368-33-7, Ch55

RL: BAC (Biological activity or effector, except adverse); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(receptor-selective retinoids as inhibitors of human head and neck squamous cell carcinoma cells)

REFERENCE COUNT: 59 THERE ARE 59 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L53 ANSWER 4 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1999:705471 HCAPLUS

DOCUMENT NUMBER: 132:89965

TITLE: Mechanism of Ubiquinol Oxidation by the bcl Complex: Different Domains of the Quinol Binding Pocket and Their Role in the Mechanism and Binding of Inhibitors
AUTHOR(S): Crofts, Antony R.; Barquera, Blanca; Gennis, Robert B.; Kuras, Richard; Guergova-Kuras, Mariana; Berry, Edward A.

CORPORATE SOURCE: Center for Biophysics and Computational Biology and Department of Biochemistry, University of Illinois at Urbana-Champaign, Urbana, IL, 61801, USA

SOURCE: Biochemistry (1999), 38(48), 15807-15826
CODEN: BICHAW; ISSN: 0006-2960

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Structures of mitochondrial ubihydroquinone:cytochrome c oxidoreductase (bcl complex) from several animal sources have provided a basis for understanding the functional mechanism at the mol. level. Using structures of the chicken complex with and without inhibitors, we analyze the effects of mutation on quinol oxidn. at the Qo site of the complex. We suggest a mechanism for the reaction that incorporates two features revealed by the structures, a movement of the iron sulfur protein between two sep. reaction domains on cytochrome c1 and cytochrome b and a bifurcated vol. for the Qo site. The vol. identified by inhibitor binding as the Qo site has two domains in which inhibitors of different classes bind differentially; a domain proximal to heme bL, where myxothiazole and .beta.-methoxyacrylate- (MOA-) type inhibitors bind (class II), and a distal domain close to the iron sulfur protein docking interface, where stigmatellin and 5-n-undecyl-6-hydroxy-4,7-dioxobenzothiaole (UHDBT) bind (class I). Displacement of one class of inhibitor by another is accounted for by the overlap of their vols., since the exit tunnel to the lipid phase forces the hydrophobic "tails" to occupy common space. We conclude that the site can contain only one "tailed" occupant, either an inhibitor or a quinol or one of their reaction products. The differential sensitivity of strains with mutations in the different domains is explained by the proximity of the affected residues to the binding domains of the inhibitors. New insights into mechanism are provided by anal. of mutations that affect changes in the ESR spectrum of the iron sulfur protein, assocd. with its interactions with the Qo-site occupant. The

structures show that all interactions with the iron sulfur protein must occur at the distal position. These include interactions between quinone, or class I inhibitors, and the reduced iron sulfur protein and formation of a reaction complex between quinol and oxidized iron sulfur protein. The step with high activation energy is after formation of the reaction complex, likely in formation of the semiquinone and subsequent disocn. of the complex into products. We suggest that further progress of the reaction requires a movement of semiquinone to the proximal position, thus mapping the bifurcated reaction to the bifurcated vol. We suggest that such a movement, together with a change in conformation of the site, would remove any semiquinone formed from further interaction with the oxidized [2Fe-2S] center and also from reaction with O₂ to form superoxide anion. We also identify two sep. reaction paths for exit of the two protons released in quinol oxidn.

IT 103455-29-4D, complexes with ubiquinol-cytochrome c reductase
RL: PRP (Properties)

(different domains of the quinol binding pocket of the bcl complex and their role in the mechanism and binding of inhibitors)

REFERENCE COUNT: 67 THERE ARE 67 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L53 ANSWER 5 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1999:674667 HCAPLUS

DOCUMENT NUMBER: 132:60924

TITLE: Effect of Inhibitors on the Ubiquinone Binding Capacity of the Primary Energy Conversion Site in the Rhodobacter capsulatus Cytochrome bcl Complex

AUTHOR(S): Sharp, R. Eryl; Gibney, Brian R.; Palmitessa, Aimee; White, Jennifer L.; Dixon, Jennifer A.; Moser, Christopher C.; Daldal, Fevzi; Dutton, P. Leslie

CORPORATE SOURCE: Johnson Research Foundation Department of Biochemistry and Biophysics and Plant Science Institute, University of Pennsylvania, Philadelphia, PA, 19104, USA

SOURCE: Biochemistry (1999), 38(45), 14973-14980

CODEN: BICHAW; ISSN: 0006-2960

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A key issue concerning the primary conversion (QO) site function in the cytochrome bcl complex is the stoichiometry of ubiquinone/ubihydroquinone occupancy. Previous evidence suggests that the QO site is able to accommodate two ubiquinone mols., the double occupancy model [Ding, H., Robertson, D. E., Daldal, F., and Dutton, P. L. (1992) Biochem. 31, 3144-3158]. In the recently reported crystal structures of the cytochrome bcl complex, no electron d. was identified in the QO site that could be ascribed to ubiquinone. To provide further insight into this issue, we have manipulated the cytochrome bcl complex QO site occupancy in photosynthetic membranes from Rhodobacter capsulatus by using inhibitor titrns. and ubiquinone extn. to modulate the amt. of ubiquinone bound in the site. The nature of the QO site occupants was probed via the sensitivity of the reduced [2Fe-2S] cluster ESR (EPR) spectra to modulation of QO site occupancy. Diphenylamine (DPA) and methoxyacrylate (MOA)-stilbene are known QO site inhibitors of the cytochrome bcl complex. Addn. of stoichiometric concns. of MOA-stilbene or excess DPA to cytochrome bcl complexes with natural levels of ubiquinone elicits the same change in the [2Fe-2S] cluster EPR spectra; the gx resonance broadens and shifts from 1.800 to 1.783. This is exactly the same signal as that obtained when there is only one ubiquinone present in the QO site.

Furthermore, addn. of MOA-stilbene or DPA to the cytochrome bcl complex depleted of ubiquinone does not alter the [2Fe-2S] cluster EPR spectral line shapes, which remain indicative of one ubiquinone or zero ubiquinones in the QO site, with broad gx resonances at 1.783 or 1.765, resp. The results are quite consistent with the QO site double occupancy model, in which MOA-stilbene and DPA inhibit by displacing one, but not both, of the QO site ubiquinones.

IT 103455-29-4

RL: BAC (Biological activity or effector, except adverse); BIOL (Biological study)

(ESR studies of the effect of **inhibitors** on ubiquinone **binding** capacity of primary energy conversion site in Rhodobacter capsulatus cytochrome bcl complex)

REFERENCE COUNT: 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L53 ANSWER 6 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1998:460644 HCAPLUS

DOCUMENT NUMBER: 129:185980

TITLE: Inhibitor binding changes domain mobility in the iron-sulfur protein of the mitochondrial bcl complex from bovine heart

AUTHOR(S): Kim, Hoeon; Xia, Di; Yu, Chang-An; Xia, Jia-Zhi; Kachurin, Anatoly M.; Zhangs, Li; Yu, Linda; Deisenhofer, Johann

CORPORATE SOURCE: Howard Hughes Medical Institute and Department of Biochemistry, University of Texas Southwestern Medical Center, Dallas, TX, 75235-9050, USA

SOURCE: Proc. Natl. Acad. Sci. U. S. A. (1998), 95(14), 8026-8033

CODEN: PNASA6; ISSN: 0027-8424

PUBLISHER: National Academy of Sciences

DOCUMENT TYPE: Journal

LANGUAGE: English

AB We have analyzed crystal structures of cytochrome bcl complexes with electron transfer inhibitors bound to the ubiquinone binding pockets Qi and/or Qo in the cytochrome b subunit. The presence or absence of the Qi inhibitor antimycin A did not affect the binding of the Qo inhibitors. Different subtypes of Qo inhibitors had dramatically different effects on the mobility of the extramembrane domain of the iron-sulfur protein (ISP): binding of 5-undecyl-6-hydroxy-4,7-dioxobenzothiazol and stigmatellin (subtype Qo-II and Qo-III, resp.) led to a fixation of the ISP domain on the surface of cytochrome b, whereas binding of myxothiazol and methoxyacrylate-stilbene (subtype Qo-I) favored release of this domain. The native structure has an empty Qo pocket and is intermediate between these extremes. On the basis of these observations we propose a model of quinone oxidn. in the bcl complex which incorporates fixed and loose states of the ISP as features important for electron transfer and, possibly, also proton transport.

IT 103455-29-4

RL: BAC (Biological activity or effector, except adverse); BIOL (Biological study)

(**inhibitor binding**-mediated changes in domain mobility in the iron-sulfur protein of the mitochondrial bcl complex from bovine heart have mechanistic implications)

L53 ANSWER 7 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1997:306198 HCAPLUS

DOCUMENT NUMBER: 127:209

TITLE: Induction of the oxidative catabolism of retinoic acid in MCF-7 cells
 AUTHOR(S): Krekels, M.D.W.G.; Verhoeven, A.; Van Dun, J.; Cools, W.; Van Hove, C.; Dillen, L.; Coene, M-C.; Wouters, W.
 CORPORATE SOURCE: Janssen Research Foundation, Department of Oncology, Beerse, B-2340, Belg.
 SOURCE: Br. J. Cancer (1997), 75(8), 1098-1104
 CODEN: BJCAAI; ISSN: 0007-0920
 PUBLISHER: Churchill Livingstone
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Cytochrome P 450-dependent oxidn. is a pathway for all-trans-retinoic acid (all-trans-RA) catabolism. Induction of this catabolic pathway was studied in MCF-7 breast cancer cells. MCF-7 cells showed low constitutive all-trans-RA catabolism. Conc.-dependent induction was obtained by preincubation of the cells with all-trans-RA (10^{-9} to 10^{-6} M). Onset of induction was fast, being detectable within 60 min, with maximal induction (45-fold) obtained after 16 h. Enzymic characterization of induced all-trans-RA catabolism showed an estd. Km value (Michaelis-Menten const.) of 0.33 μ M and a Vmax value (maximal velocity of an enzyme-catalyzed reaction) of 54.5 fmol polar all-trans-RA metabolites 106 cells $^{-1}$ h $^{-1}$. These kinetic parameters represent the overall formation of polar metabolites from all-trans-RA. Induction of all-trans-RA catabolism was also obtained with other retinoids, CH55 μ chgt. 13-cis-RA = All-trans-RA > 9-cis-RA > 4-keto-all-trans-RA > 4-keto-13-cis-RA > retinol. The potency of the retinoids to induce all-trans-RA catabolism was correlated to their retinoic acid receptor affinity (Crettaz et al, 1990; Repa et al, 1990; Sani et al, 1990). Induction of all-trans-RA catabolism was inhibited by actinomycin D. Furthermore, all-trans-RA did not increase cytosolic retinoic acid-binding protein (CRABP) mRNA levels. These data suggest that induction of all-trans-RA catabolism in MCF-7 cells is a retinoic acid receptor-mediated gene transcriptional event. Induced all-trans-RA catabolism was inhibited by various retinoids with decreasing potency in the order: all-trans-RA > 4-keto-all-trans-RA > 13-cis-RA > 9-cis-RA > 4-keto-13-cis-RA > retinol > CH55. The antitumoral compd. liarozole-fumarate inhibited all-trans-RA catabolism with a potency similar to that of all-trans-RA.

IT 110368-33-7, CH55

RL: BAC (Biological activity or effector, except adverse); BIOL.
 (Biological study)
 (induction of oxidative catabolism of retinoic acid in MCF-7 cells by retinoids by **receptor** interaction and **inhibition** by liarozole and actinomycin D in relation to retinoic acid-binding protein)

L53 ANSWER 8 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1997:143330 HCAPLUS
 DOCUMENT NUMBER: 126:246352
 TITLE: Inhibition of IL-1-induced IL-6 production by synthetic retinoids
 AUTHOR(S): Kagechika, Hiroyuki; Kawachi, Emiko; Fukasawa, Hiroshi; Saito, Go; Iwanami, Naoko; Umemiya, Hiroki; Hashimoto, Yuichi; Shudo, Koichi
 CORPORATE SOURCE: Faculty of Pharmaceutical Sciences, University of Tokyo, Tokyo, 113, Japan
 SOURCE: Biochem. Biophys. Res. Commun. (1997), 231(2), 243-248
 CODEN: BBRCA9; ISSN: 0006-291X
 PUBLISHER: Academic
 DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effects of retinoids and retinoid antagonists on IL-6 prodn. in MC3T3-E1 cells were investigated. None of the synthetic retinoids examd. stimulated IL-6 prodn., but all of them strongly inhibited IL-6 prodn. induced by mouse IL-1.alpha.. Their inhibitory activities correlated well with their differentiation-inducing activities in HL-60 assay or their binding affinities to nuclear retinoic acid receptors (RARs). Among three retinoid antagonists, two weak antagonists exhibited similar inhibition of mouse IL-1.alpha.-induced IL-6 prodn., whereas a potent retinoid antagonist, 4-(13H-10,11,12,13-tetrahydro-10,10,13,13,15-pentamethyl-dinaphtho[2,3-b][1,2-e]diazepin-7-yl)benzoic acid (LE540), enhanced IL-6 prodn. under the same conditions.

IT 110368-33-7, Ch55

RL: BAC (Biological activity or effector, except adverse); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (inhibition of IL-1-induced IL-6 prodn. by synthetic retinoids and retinoid antagonists in relation to differentiation-inducing activity and retinoid **receptor binding** and structure)

L53 ANSWER 9 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1996:135308 HCAPLUS

DOCUMENT NUMBER: 124:221563

TITLE: The RAR-RXR as well as the RXR-RXR pathway is involved in signaling growth inhibition of human CD34+ erythroid progenitor cells

AUTHOR(S): Rusten, Leiv S.; Dybedal, Ingunn; Blomhoff, Heidi Kiil; Blomhoff, Rune; Smeland, Erlend B.; Jacobsen, Sten Eirik W.

CORPORATE SOURCE: Dep. Immunol., Institute Cancer Research, Norwegian Radium Hospital, Oslo, N-0310, Norway

SOURCE: Blood (1996), 87(5), 1728-36
CODEN: BLOOAW; ISSN: 0006-4971

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Previous studies have shown that retinoic acid (RA), similar to tumor necrosis factor-.alpha. (TNF-.alpha.), can act as a bifunctional regulator of the growth of bone marrow progenitors, in that it can stimulate granulocyte-macrophage colony-stimulating factor (GM-CSF)- or interleukin-3 (IL-3)-induced GM colony formation, but potently inhibit G-CSF-induced growth. The present study, using highly enriched human CD34+ as well as Lin- murine bone marrow progenitor cells, demonstrates a potent inhibitory effect of 9-cis-RA on burst-forming unit-erythroid (BFU-e) colony formation regardless of the cytokine stimulating growth. Specifically, 9-cis-RA potently inhibited the growth of BFU-E in response to erythropoietin (Epo) (100%), stem cell factor (SCF) + Epo (92%), IL-3 + Epo (97%), IL-4 + Epo (88%), and IL-9 + Epo (100%). Erythroid colony growth was also inhibited when CD34+ progenitors were seeded at one cell per well, suggesting a direct action of RA. Using synthetic ligands to retinoic acid receptors (RARs) and retinoid X receptors (RXRs) that selectively bind and activate RAR-RXR or RXR-RXR dimers, resp., the authors dissected the involvement of the two retinoid response pathways in the regulation of normal myeloid and erythroid progenitor cell growth. Transactivation studies showed that both the RAR (Ro 13-7410) and RXR (Ro 25-6603 and Ro 25-7386) ligands were highly selective at 100 nmol/L. At this concn., Ro 13-7410 potently inhibited G-CSF-stimulated myeloid as well as SCF + Epo-induced erythroid colony growth. At the same concn., Ro 25-6603 and Ro 25-7386 had little or no effect on G-CSF-induced colony formation, whereas they inhibited 75% and 53%, resp., of SCF +

Epo-stimulated BFU-E colony growth. Thus, the RAR-RXR response pathway can signal growth inhibition of normal bone marrow myeloid and erythroid progenitor cells. In addn., the authors demonstrate a unique involvement of the RXR-RXR pathway in mediating growth inhibition of erythroid but not myeloid progenitor cells.

IT 173792-73-9, Ro 25-6603

RL: BAC (Biological activity or effector, except adverse); BIOL (Biological study)

(RAR-RXR and RXR-RXR **receptor** pathway signaling of cytokine-stimulated growth **inhibition** of human CD34+ erythroid progenitor cells)

L53 ANSWER 10 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1995:836520 HCAPLUS

DOCUMENT NUMBER: 124:209

TITLE: Correlation of retinoid binding affinity to retinoic acid receptor .alpha. with retinoid inhibition of growth of estrogen receptor-positive MCF-7 mammary carcinoma cells

AUTHOR(S): Dawson, Marcia I.; Chao, Wan-ru; Pine, Polly; Jong, Ling; Hobbs, Peter D.; Rudd, Colette K.; Quick, Timothy C.; Niles, Richard M.; Zhang, Xiao-kun; et al.

CORPORATE SOURCE: Life Sciences Div., SRI Internatl., Menlo Park, CA, 94025, USA

SOURCE: Cancer Res. (1995), 55(19), 4446-51

CODEN: CNREA8; ISSN: 0008-5472

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Both anchorage-dependent growth and anchorage-independent growth of the estrogen receptor-pos. mammary carcinoma cell line MCF-7 are inhibited by all-trans-retinoic acid. This cell line has nuclear retinoic acid receptors (RARs) .alpha. and .gamma.. The natural retinoids all-trans-retinoic acid and 9-cis-retinoic acid and a series of 12 conformationally restricted retinoids, which showed a range of binding selectivities for these receptors and had either agonist or antagonist activity for gene transcriptional activation by the RARs, were evaluated for their abilities to inhibit anchorage-dependent (adherent) and anchorage-independent (clonal) growth of MCF-7 cells. Correlation analyses were performed to relate growth inhibition by these retinoids with their binding affinity to RAR.alpha. or RAR.gamma.. Inhibition of anchorage-dependent growth in culture after 7 days of retinoid treatment correlated with binding to RAR.alpha. and not to RAR.gamma.. Both the RAR.alpha.-selective retinoid agonists and the two RAR antagonists that were evaluated inhibited adherent cell growth. The RAR.gamma.-selective agonists had very low growth inhibitory activity (<10%) at concns. as high as 12.5 .mu.M. These results suggest that RAR.alpha. is the retinoid receptor involved in the inhibition of adherent cell growth by retinoids and that transcriptional activation by this receptor on a RAR response element does not appear to be required for this process to occur. For this series of retinoids, inhibition of anchorage-independent growth after 21 days of retinoid treatment only correlated with binding affinity to RAR.alpha. for the retinoid agonists, although the RAR.gamma.-selective retinoids displayed weak activity. The RAR antagonists were very poor inhibitors of growth. These results suggest that activation of gene transcription by RAR.alpha. appears to be required for inhibition of anchorage-independent growth by retinoids in this estrogen receptor-pos. mammary carcinoma cell line.

IT 75664-66-3, SR 3983 138254-19-0, SR 3986

RL: BAC (Biological activity or effector, except adverse); BPR (Biological

process); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)

(correlation of retinoid **binding** affinity to retinoic acid **receptor** .alpha. with retinoid **inhibition** of growth of estrogen **receptor**-pos. MCF-7 mammary carcinoma cells in relation to structure)

L53 ANSWER 11 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1994:217212 HCAPLUS

DOCUMENT NUMBER: 120:217212

TITLE: 6,6-Disubstituted hex-5-enoic acid derivatives as combined thromboxane A2 receptor antagonists and synthetase inhibitors

AUTHOR(S): Söyka, Rainer; Heckel, Armin; Nickl, Josef; Eisert, Wolfgang; Mueller, Thomas H.; Weisenberger, Hans
CORPORATE SOURCE: Res. Dep., Dr. Karl Thomae GmbH, Biberach, 88397, Germany


SOURCE: J. Med. Chem. (1994), 37(1), 26-39

CODEN: JMCMAR; ISSN: 0022-2623

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



AB A series of .omega.-disubstituted alkenoic acid derivs. were design and synthesized as antithrombotic inhibitors of thromboxane A2 synthetase and thromboxane A2 receptor antagonists. Hexenoic acid derivs. with a 3-pyridyl group and a 4-(2-benzenesulfonamidoethyl)phenyl substituent were found to be optimal with regard to the dual mode of action. The most potent compd., (E)-6-(4-(2-(((4-chlorophenyl)sulfonyl)amino)ethyl)phenyl)-6-(3-pyridyl)hex-5-enoic acid (I), inhibits thromboxane A2 synthetase in gel-filtered human platelets with an IC50 value of 4.5 +/- 0.5 nM, whereas an inhibitory effect on cyclooxygenase is seen only at a much higher concn. (IC50; 240 .mu.M). Radioligand-binding studies with [3H]SQ 29,548 in washed human platelets revealed that I blocks the prostaglandin H2/thromboxane A2 receptor with an IC50 fo 19 +/- 5 nM (n = 5) and is therefore 85-fold more potent than another combined thromboxane A2 synthetase inhibitor/receptor antagonist, Ridogrel. I inhibits the collagen-induced platelet aggregation in human platelet-rich plasma and whole blood with an EC50 of 1 .mu.M (Ridogrel: 16 .mu.M) and 100 nM, resp., and was selected for further development.

IT 153732-10-6P 153732-27-5P

RL: SPN (Synthetic preparation); PREP (Preparation)

(prepn. and thromboxane A2 **receptor** antagonist and synthetase
inhibitor activity of)

L53 ANSWER 12 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1994:182467 HCAPLUS

DOCUMENT NUMBER: 120:182467

TITLE: Correlation of the ability of retinoids to inhibit
promoter-induced anchorage-independent growth of JB6
mouse epidermal cells with their activation of
retinoic acid receptor .gamma.

AUTHOR(S): Rudd, Colette J.; Mansbridge, Jonathan N.; Suing,
Kathryn D.; Dawson, Marcia I.

CORPORATE SOURCE: Life Sci. Div., SRI Int., Menlo Park, CA, 94025, USA

SOURCE: Cancer Lett. (Shannon, Irel.) (1993), 73(1), 41-9

CODEN: CALEDQ; ISSN: 0304-3835

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Retinoids inhibit the biol. effects induced in mouse epidermal cells by
the tumor promoter 12-O-tetradecanoyl-phorbol-13-acetate (TPA). Specific
nuclear retinoic acid receptors (RARs) have been identified in the
epidermis, but the specific receptor that mediates the inhibitory response
by retinoids is not established. Retinoic acid and six conformationally
restricted retinoids were evaluated in an in vitro bioassay using the JB6
mouse epidermal cell line. These activities were then compared with the
ability of these retinoids to activate the RARs in transient transfection
assays for transcriptional activation to identify the retinoid receptor
involved in inhibiting TPA-induced anchorage-independent growth. The
retinoids inhibited TPA-induced colony formation of JB6 cells in semisolid
medium at concns. that were not toxic based on colony formation of
attached cells. These concns. ranged from less than 10⁻⁹-10⁻⁶ M.
4-(5,6,7,8-Tetrahydro-5,5,8,8-tetramethylanthracen-2-yl)benzoic acid
(TTAB) was the most potent retinoid, with an EC50 of 0.8 nM. Both
RAR.alpha. and RAR.gamma. were expressed in JB6 cells. Expression of
RAR.beta. was not detected in these cells using a polymerase chain
reaction assay, consistent with its extremely low level in mouse skin.
Inhibition of the TPA response by these retinoids in JB6 cells correlated
only with their transcriptional activation of RAR.gamma., but not with
that of RAR.alpha.. These results suggest that RAR.gamma. is most
probably the receptor that mediates the chemopreventive effects of
retinoids in mouse epidermis.

IT 75664-66-3, SR 3983

RL: BIOL (Biological study)

(epidermal cancer **inhibition** by, retinoic acid
receptors in)

L53 ANSWER 13 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1991:511425 HCAPLUS

DOCUMENT NUMBER: 115:111425

TITLE: Expression of nuclear retinoic acid receptors in
normal tracheobronchial cells and in lung carcinoma
cells

AUTHOR(S): Nervi, Clara; Vollberg, Thomas M.; George, Margaret
D.; Zelent, Arthur; Chambon, Pierre; Jetten, Anton M.

CORPORATE SOURCE: Lab. Pulm. Pathobiol., Natl. Inst. Environ. Health
Sci., Research Triangle Park, NC, 27709, USA

SOURCE: Exp. Cell Res. (1991), 195(1), 163-70

CODEN: ECREAL; ISSN: 0014-4827

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Retinoids are important regulators of the growth and differentiation of tracheobronchial epithelial cells. To det. the mechanism of action of retinoids in these cells, the expression of nuclear retinoic acid receptors (RARs) was examd. in normal human and rabbit tracheobronchial epithelial (HBE and RbTE, resp.) cells and in several lung carcinoma cell lines. A specific nuclear RAR activity with a mol. wt. of 50,000 was identified in these cells. A correlation was found between the binding of several retinoids to this RAR and their ability to inhibit transglutaminase Type I activity. Normal HBE and RbTE cells contained 2 RAR.alpha. mRNA transcripts, 2.6 and 3.5 kb in size, and one 3.1-kb RAR.gamma. transcript. RAR.beta. transcripts were undetectable in HBE cells. RAR expression was unchanged during squamous differentiation. Treatment of HBE and RbTE cells with 100 nM retinoic acid increased RAR.beta. mRNA expression but did not change the levels of RAR.alpha. and RAR.gamma.. In contrast, retinoic acid suppressed in these cells the level of involucrin, transglutaminase Type I, and SQ37 mRNA. In comparison with normal HBE cells, certain lung carcinoma cell lines appear to have an altered expression of RAR.beta. and RAR.gamma.. Human bronchial fibroblasts (HBF) expressed RAR.alpha. and RAR.gamma. transcripts of the same size as HBE cells. HBF cells contain low levels of a 2.9- and 3.3-kb RAR.beta. mRNA. Treatment of HBF cells with retinoic acid increased the level of RAR.beta. mRNA in a time-dependent manner; the maximal induction was .apprx.15-fold. Apparently, RARs are involved in the suppression of squamous differentiation in tracheobronchial epithelial cells and lung fibroblasts are target cells for retinoids.

IT 110368-33-7, Ch55 110368-34-8, Ch 30 119567-93-0

, Ch 20 119567-94-1, Ch 40

RL: BIOL (Biological study)

(retinoic acid **receptor binding** of and squamous differentiation **inhibition** by, in tracheobronchial epithelium)

L53 ANSWER 14 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1991:422233 HCAPLUS

DOCUMENT NUMBER: 115:22233

TITLE: Thromboxane A2 receptor antagonists for the inhibition of vasospasms and thrombocyte aggregation, following angioplasty

INVENTOR(S): Ondetti, Miguel Angel; Ogletree, Martin Lawrence; Harris, Don Navarro

PATENT ASSIGNEE(S): Squibb, E. R., and Sons, Inc., USA

SOURCE: Ger. Offen., 9 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 4013680	A1	19901108	DE 1990-4013680	19900427
CA 2012852	AA	19901101	CA 1990-2012852	19900322
GB 2231795	A1	19901128	GB 1990-9636	19900430
JP 02295922	A2	19901206	JP 1990-116704	19900501
FR 2646351	A1	19901102	FR 1990-5547	19900502
PRIORITY APPLN. INFO.:			US 1989-346447	19890501

OTHER SOURCE(S): MARPAT 115:22233

GI



I

AB The 7-oxabicycloheptane prostaglandin analogs I [A = CH:CH, CH₂CH₂; B = A, C.tplbond.C; X = OH, tetrazolyl, CO₂R, CONH₂; Y = (un)substituted-alkyl, aralkyl, alkenyl, (un)substituted pyridyl, etc.; R = H, alkyl; Z = R, aryl, etc.; m = 1-8] are thromboxane A₂ receptor antagonists, useful for inhibiting vasospasms and thrombocyte aggregation following angioplasty (no data). Injection solns. comprised SQ-29548 2500, methylparaben 5, propylparaben 1, and NaCl 25 mg, in 5 L water.

IT 123048-11-3, SQ 28568

RL: BIOL (Biological study)

(thromboxane A₂ **receptor** antagonist, vasospasm and thrombocyte aggregation **inhibition** by, following angioplasty)

L53 ANSWER 15 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1991:178366 HCAPLUS

DOCUMENT NUMBER: 114:178366

TITLE: Retinoids induce tissue transglutaminase in NIH-3T3 cells

AUTHOR(S): Cai, D.; Ben, T.; De Luca, L. M.

CORPORATE SOURCE: Lab. Cell. Carcinog., Tumor Promot., Natl. Cancer Inst., Bethesda, MD, 20892, USA

SOURCE: Biochem. Biophys. Res. Commun. (1991), 175(3), 1119-24
CODEN: BBRCA9; ISSN: 0006-291X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB All-trans and 13-cis-retinoic acid (RA) as well as the synthetic compd. CH-55 enhanced tissue transglutaminase activity as they increased NIH-3T3 cell adhesiveness. The 4-hydroxyphenylretinamide with low activity in inducing attachment, lectin binding and growth inhibition failed to induce transglutaminase. Thyroxine, a compd. with a response element common to RA, was inactive. The tumor promoter 12-tetradecanoyl-phorbol-13-acetate, which increases adhesiveness with different kinetics than RA, failed to enhance transglutaminase. Thus, retinoids with biol. activity in inducing adhesion, growth inhibition and lectin binding, are also active in inducing transglutaminase activity.

IT 110368-33-7, CH-55

RL: BIOL (Biological study)

(tissue transglutaminase induced by, cell adhesion and growth **inhibition** and lectin **binding** in relation to)

L53 ANSWER 16 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1988:504166 HCAPLUS

DOCUMENT NUMBER: 109:104166

TITLE: Inhibition of ornithine decarboxylase induction by retinobenzoic acids in relation to their binding affinities to cellular retinoid-binding proteins

AUTHOR(S): Takagi, Kanji; Suganuma, Masami; Kagechika, Hiroyuki; Shudo, Koichi; Ninomiya, Mitsuo; Muto, Yasutoshi; Fujiki, Hirota

CORPORATE SOURCE: Cancer Prev. Div., Natl. Cancer Cent. Res. Inst.,
Tokyo, 104, Japan
SOURCE: J. Cancer Res. Clin. Oncol. (1988), 114(3), 221-4
CODEN: JCROD7; ISSN: 0171-5216

DOCUMENT TYPE: Journal
LANGUAGE: English

AB Retinobenzoic acids induce differentiation of human promyelocytic leukemia cells (HL-60). Like retinoic acid, 14 retinobenzoic acids inhibited the induction of ornithine decarboxylase (ODC) by teleocidin in mouse skin. The mechanism(s) of inhibition of ODC induction by 7 retinobenzoic acids, Am 80, Am 81, Am 580, Am 590, Am 68, Sa 80, and Ch 55 was compared with those for all-trans-retinoic acid and the arotinoid compd. 19. Application of 114 nmol of Am 80, Am 81, Am 580, Am 590, Am 68, Sa 80, or Ch 55, 10 min before 11.4 nmol of teleocidin, resulted in 76.7, 82.0, 76.2, 28.3, 48.4, 58.6, and 85.1% inhibition of ODC induction, resp. Since all-trans-retinoic acid and compd. 19 were also inhibitory, the authors detd. whether retinobenzoic acids bind to cellular retinoic acid-binding protein (CRABP) isolated from bovine adrenal glands. Am 80 and Am 580 inhibited the specific binding of [3H]retinoic acid to CRABP, but also showed less affinity than authentic unlabeled retinoic acid and compd. 19. Am 81, Am 590, Am 68, Sa 80, and Ch 55 at up to 10 .mu.M were not effective competitors of the binding of either [3H]retinoic acid or [3H]retinol. Apparently, the inhibition of ODC induction can be mediated by pathways that do not involve CRABP or the cellular retinol-binding protein.

IT 110368-33-7, Ch 55

RL: BIOL (Biological study)
(ornithine decarboxylase inhibition by; cellular retinoid-binding protein and structure in relation to)

L53 ANSWER 17 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1988:469330 HCAPLUS

DOCUMENT NUMBER: 109:69330

TITLE: Characterization of binding of the methoxyacrylate inhibitors to mitochondrial cytochrome c reductase

AUTHOR(S): Brandt, Ulrich; Schaegger, Hermann; Von Jagow, Gebhard
CORPORATE SOURCE: Inst. Phys. Biochem., Univ. Muenchen, Munich, Fed. Rep. Ger.

SOURCE: Eur. J. Biochem. (1988), 173(3), 499-506
CODEN: EJBCAI; ISSN: 0014-2956

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The 3 (E)-.beta.-methoxyacrylate (MOA) inhibitors oudemansin A, strobilurin A, and MOA stilbene, which differ by >1 order of magnitude in their binding affinity to mitochondrial ubihydroquinone-cytochrome c oxidoreductase (bc1 complex), bind to a site that is not identical to the binding site for ubihydroquinone, the substrate of the outer ubiquinone reaction site (Q0 center). Although the ubihydroquinone mol. is still bound in the presence of the MOA inhibitors, its electrons cannot be transferred to the Fe-S center. A shift of the relative position of the ubihydroquinone mol. in the reaction center due to a conformational distortion of cytochrome b induced by the binding of the MOA inhibitor seems to be the reason for the blocked electron transfer. Ubihydroquinone affects the dissocn. const. values of all 3 MOA inhibitors tested; the values are raised by a const. factor of 2, although the inhibitors bind with quite different affinity. The Fe-S protein is not involved in the binding of the MOA inhibitors. These results have direct implications for the proper use of MOA inhibitors in expts. designed to analyze the structure/mechanism relationship in cytochrome c reductase. In

particular, point mutations recently described in MOA inhibitor-resistant mutants can no longer be taken to affect necessarily the ubihydroquinone-binding site.

IT 103455-29-4

RL: BIOL (Biological study)
(ubiquinol-cytochrome c reductase **inhibition** by, kinetics of, **binding** site in)

L53 ANSWER 18 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1987:526601 HCAPLUS

DOCUMENT NUMBER: 107:126601

TITLE: New benzoic acid derivatives with retinoid activity:
~~lack of direct correlation between biological activity~~
and binding to cellular retinoic acid binding protein

AUTHOR(S): Jetten, Anton M.; Anderson, K.; Deas, M. A.;
Kagechika, H.; Lotan, R.; Rearick, J. I.; Shudo, K.

CORPORATE SOURCE: Lab. Pulm. Pathobiol., Natl. Inst. Environ. Health
Sci., Research Triangle Park, NC, 27709, USA

SOURCE: Cancer Res. (1987), 47(13), 3523-7

CODEN: CNREA8; ISSN: 0008-5472

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The biol. activity of several newly synthesized benzoic acid derivs. of the Am- and Ch- series, which are structurally different from retinoic acid and arotinoids, was examd. These compds. inhibit squamous cell differentiation of rabbit tracheal epithelial cells in vitro as indicated by the inhibition of transglutaminase Type I and cholesterol 3-sulfate levels. In contrast to the inhibition of differentiation in rabbit tracheal cells, these compds. induce differentiation of mouse embryonal carcinoma F9 and human promyelocytic leukemia HL60 cells. The Am- and Ch- series of compds. also affect several parameters of cell proliferation. These agents are very potent inhibitors of growth of melanoma S91 cells and inhibit the induction of ornithine decarboxylase activity by phorbol 12-myristate 13-acetate in 3T6 fibroblasts. These results show that the Am- and Ch- derivs. elicit in several cell systems the same cellular responses as retinoic acid. Apparently, they exhibit mechanism(s) of action similar to those of retinoids. Comparison of the biol. response with the binding capacity to the cellular retinoic acid-binding protein shows a lack of a direct correlation.

IT 110368-33-7 110368-34-8

RL: BIOL (Biological study)
(cell differentiation **modulation** and growth
inhibition by, **binding** to cellular retinoic acid-
binding protein correlation with)

L53 ANSWER 19 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1987:78495 HCAPLUS

DOCUMENT NUMBER: 106:78495

TITLE: Quantitation of drug levels and platelet receptor
blockade caused by a thromboxane antagonist

AUTHOR(S): Friedhoff, Lawrence T.; Manning, J.; Funke, P. T.;
Ivashkiv, E.; Tu, J.; Cooper, W.; Willard, D. A.

CORPORATE SOURCE: Squibb Inst. Med. Res., Princeton, NJ, USA

SOURCE: Clin. Pharmacol. Ther. (St. Louis) (1986), 40(6),
634-42

CODEN: CLPTAT; ISSN: 0009-9236

DOCUMENT TYPE: Journal

LANGUAGE: English

AB SQ 28,668 (I) [87983-63-9] is a structural analog of

thromboxane A2 [57576-52-0]. It **inhibits** the effects of thromboxane in vitro. Healthy male subjects were given either placebo or 3 equal daily doses of I ranging from 25 to 1200 mg. Plasma drug concns. increased in a dose-dependent manner. The shape of the plasma drug concn.-time curve was consistent with enterohepatic recirculation. The effects of I on ex vivo platelet aggregation suggested that I is a specific competitive antagonist of thromboxane A2 with a platelet **receptor** dissocn. const. (estd. by Schild anal.) of .apprx.19 nM. Approx. 94% occupation of thromboxane **receptors** by I was required to produce a small but measurable increase of the template bleeding time. Dose-ranging studies of antithrombotic drugs are difficult and expensive. For this reason, a method was developed that allows estn. of the dose of a thromboxane **receptor** antagonist that would be expected to be therapeutically equiv. to a given dose of aspirin.

IT 87983-63-9

RL: BIOL (Biological study)
(platelet aggregation and thromboxane A2 **receptors**
inhibition by, in humans)

L53 ANSWER 20 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1986:230538 HCAPLUS

DOCUMENT NUMBER: 104:230538

TITLE: t-PA composition and getting it into the blood stream

INVENTOR(S): Sarnoff, Stanley Jay

PATENT ASSIGNEE(S): Survival Technology, Inc., USA

SOURCE: PCT Int. Appl., 42 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 5

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 8601104	A1	19860227	WO 1985-US1503	19850808
W: AU, BR, JP				
RW: AT, BE, CH, DE, FR, GB, IT, LU, NL, SE				
US 4661469	A	19870428	US 1985-708845	19850306
AU 8546744	A1	19860307	AU 1985-46744	19850808
AU 589135	B2	19891005		
EP 191056	A1	19860820	EP 1985-903966	19850808
R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE				
JP 61502892	T2	19861211	JP 1985-503532	19850808
IL 76120	A1	19901223	IL 1985-76120	19850818
WO 8605095	A1	19860912	WO 1986-US473	19860306
W: DK, FI, HU, KR, NO				
ZA 8601695	A	19861029	ZA 1986-1695	19860306
US 5002930	A	19910326	US 1987-19564	19870227
US 32919	E	19890509	US 1987-66732	19870626
AU 9053572	A1	19911030	AU 1990-53572	19900330
IL 93973	A1	19961031	IL 1990-93973	19900401

PRIORITY APPLN. INFO.:

US 1984-638695	19840808
US 1985-708845	19850306
US 1985-716705	19850327
WO 1985-US1503	19850808
US 1985-782441	19851001
WO 1990-US1633	19900330

AB Tissue plasminogen activator (t-PA) absorption i.m. is enhanced by simultaneous application of hydroxylamine or a nontoxic salt thereof, and

optionally by elec. stimulation at the injection site to increase muscle blood flow, addn. of platelet aggregation inhibitors, etc. Thus, injection (i.m.) of t-PA into rabbits along with hydroxylamine.HCl increased t-PA blood plasma levels 40-fold as compared with t-PA injection alone (measured 5 min after injection).

IT 93060-40-3

RL: BIOL (Biological study)
(thromboxane A **receptor inhibition** by, after
plasminogen activator application)

L53 ANSWER 21 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1985:108185 HCAPLUS

DOCUMENT NUMBER: 102:108185

TITLE: Efficacy of Bayticol (flumethrin) against ixodid ticks under field conditions, and its tolerance in cattle

AUTHOR(S): Hamel, H. D.; Dorn, H.

CORPORATE SOURCE: PH-Vet. Ber., Bayer A.-G., Cologne, 5000/80, Fed. Rep. Ger.

SOURCE: Acarol., [Proc. Int. Congr. Acarol.], 6th (1984), Meeting Date 1982, Volume 2, 1263-8. Editor(s): Griffiths, Donald Alister; Bowman, Clive Edward. Horwood: Chichester, UK.

CODEN: 52JKAI

DOCUMENT TYPE: Conference

LANGUAGE: English

AB Field trials have been conducted in various tick-enzootic countries, in order to evaluate the efficacy of flumethrin [69770-45-2] against one-host and multi-host tick species. A range of concns. was tested to det. the optimum use rates for plunge dipping and spray-race treatments. Use rates of 30 ppm against one-host ticks and 40 ppm against multi-host ticks such as Hyalomma truncatum, could be recommended, based on the results of oviposition **inhibition** tests in the lab. A prolonged protective period of >7 days was found, which correlated with the larvicidal activity of flumethrin as evaluated in in vitro tests. Flumethrin was well tolerated by cattle and other domestic animals without any side effects after a dermal application of concns. >30-50 times the recommended use rate. No detectable residues were present in milk and edible tissues after spray treatments of .ltoreq.200 ppm.

IT 69770-45-2

RL: BIOL (Biological study)
(ixodid **control** by, on cattle)

L53 ANSWER 22 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1985:108184 HCAPLUS

DOCUMENT NUMBER: 102:108184

TITLE: Biological evaluation of flumethrin, a new synthetic pyrethroid for the control of ticks

AUTHOR(S): Stendel, W.; Fuchs, R.

CORPORATE SOURCE: Inst. Chemother., Bayer A.-G., Wuppertal, D-5600/1, Fed. Rep. Ger.

SOURCE: Acarol., [Proc. Int. Congr. Acarol.], 6th (1984), Meeting Date 1982, Volume 2, 1252-62. Editor(s): Griffiths, Donald Alister; Bowman, Clive Edward. Horwood: Chichester, UK.

CODEN: 52JKAI

DOCUMENT TYPE: Conference

LANGUAGE: English

AB Flumethrin (I) [69770-45-2] (0.5-120 ppm) **inhibited** in vitro the deposition of fertile eggs by fully-engorged females of

Boophilus microplus, B. decoloratus, and B. annulatus, including organophosphorus insecticide-resistant strains. Similar results were shown on multi-host Rhipicephalus and Amblyomma species. Larvicidal activity was shown by 0.01-1 ppm I, with little difference between species. On cattle, 30 ppm I was effective against Boophilus, and 40 ppm I against the multi-host ticks.

IT 69770-45-2

RL: BIOL (Biological study)
(tick control by, on cattle)

L53 ANSWER 23 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1985:1050 HCAPLUS

DOCUMENT NUMBER: 102:1050

TITLE: Effects of SQ 27,427, a thromboxane A2 receptor antagonist, in the human platelet and isolated smooth muscle

AUTHOR(S): Harris, Don N.; Greenberg, Roland; Phillips, Marie B.; Michel, Inge M.; Goldenberg, Harold J.; Haslanger, Martin F.; Steinbacher, Thomas E.

CORPORATE SOURCE: Squibb Inst. Med. Res., Princeton, NJ, 08540, USA


SOURCE: Eur. J. Pharmacol. (1984), 103(1-2), 9-18

CODEN: EJPHAZ; ISSN: 0014-2999

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



I

AB The TXA2 [57576-52-0] **receptor** antagonist properties of SQ 27427 (I) [93060-40-3] were studied in vitro, both in the human platelet and various isolated smooth muscle preps. I was a potent **inhibitor** of human platelet aggregation induced by arachidonic acid [506-32-1], ADP [58-64-0], epinephrine [51-43-4], collagen, and the stable TXA2 agonists 9,11-azoPGH2 [57712-08-0] and SQ 26655 [82337-14-2]. **Inhibition** of platelet aggregation was achieved at I concns. that did not alter TXB2 levels. I weakly **inhibited** the formation of TXB2 from arachidonic acid and had no effect on the synthesis of PGE2 or PGI2 from arachidonic acid. I was also a weak stimulator of platelet adenylate cyclase [9012-42-4], being only 0.1% as potent as PGI2. In isolated smooth muscle expts., I was a potent and specific TXA2 **receptor** antagonist. It caused competitive antagonism of 9,11-azoPGH2-induced contractions of vascular, respiratory, and gastrointestinal smooth muscles. This antagonism was specific, as responses to norepinephrine, serotonin, PGE2, PGI2, PGF2.alpha., histamine, carbachol, and KCl were not altered by I.

IT 93060-40-3

RL: BAC (Biological activity or effector, except adverse); BIOL (Biological study)
(as TXA2 **receptor** antagonist)

L53 ANSWER 24 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1984:604872 HCAPLUS

DOCUMENT NUMBER: 101:204872

TITLE: Thromboxane receptor antagonist properties of SQ 27,427 in the anesthetized guinea pig

AUTHOR(S): Greenberg, Roland; Steinbacher, Thomas E.; Harris, Don N.; Haslanger, Martin F.

CORPORATE SOURCE: Squibb Inst. Med. Res., Princeton, NJ, 08540, USA

SOURCE: Eur. J. Pharmacol. (1984), 103(1-2), 19-24

CODEN: EJPHAZ; ISSN: 0014-2999

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The TXA2 [57576-52-0] **receptor** antagonist properties of SQ 27427 [92843-52-2], were studied in vivo in the anesthetized guinea pig where changes in pulmonary resistance, dynamic compliance, and mean arterial blood pressure were measured. Both the bronchoconstrictor and pressor responses to arachidonic acid (AA) [506-32-1] and to the stable TXA2 mimic 9,11-azoPGH2 (AZO) [57712-08-0] were taken as indexes of in vivo TXA2 **receptor** activation. The administration of SQ 27427 (0.1-1.0 mg/kg, i.v.; 10.0 mg/kg, p.o.) caused dose-related **inhibitions** of both AA- and AZO-induced bronchoconstriction. Relative specificity of this antagonism was evidenced by the failure of SQ 27427 (1.0 mg/kg, i.v.) to **inhibit** histamine-induced bronchoconstriction. In the same expts. the pressor response to AA was reversed to a depressor response by SQ 27427. This reversal was abolished by indomethacin. The pressor response to AZO was antagonized by SQ 27427, but not by indomethacin. The reversal of the pressor response to AA by SQ 27427 may be due to the unmasking of the depressor effect of a cyclooxygenase product, i.e., prostacyclin. Thus, SQ 27427 is a relatively specific TXA2 **receptor** antagonist in vivo in the guinea pig.

IT 92843-52-2

RL: BIOL (Biological study)
(as TXA2 **receptor** antagonist)

L53 ANSWER 25 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1984:137789 HCAPLUS

DOCUMENT NUMBER: 100:137789

TITLE: Relationship between binding affinities to cellular retinoic acid-binding protein and biological potency of a new series of retinoids

AUTHOR(S): Sani, Brahma P.; Dawson, Marcia I.; Hobbs, Peter D.; Chan, Rebecca L. S.; Schiff, Leonard J.

CORPORATE SOURCE: Kettering Meyer Lab., South. Res. Inst., Birmingham, AL, 35255-5305, USA

SOURCE: Cancer Res. (1984), 44(1), 190-5

CODEN: CNREA8; ISSN: 0008-5472

DOCUMENT TYPE: Journal

LANGUAGE: English

AB **Binding** affinities of a new and unusual series of retinoic acid [302-79-4] analogs to cellular retinoic acid-**binding** protein, a possible mediator of their biol. function in the **control** of differentiation and tumorigenesis, and to serum albumin, their plasma transport protein, were detd. Also, biol. activity of these retinoids in the reversal of keratinization in hamster tracheal organ cultures was assessed and compared with their **binding** affinities. Analogs that possessed high biol. activity showed high **binding** efficiency to cellular retinoic acid-**binding** protein. Those

that were biol. less active were poor **binders** to the **binding** protein. Three retinoids, 4657-57 [75664-66-3], 3920-59 [86238-80-4], and 4445-75 [86471-13-8], which showed 90-100% **binding** efficiency of that of retinoic acid for cellular retinoic acid-**binding** protein, expressed high biol. activity detectable in the range of 10-10M as against 10-11M for retinoic acid. The correlation noticed in these 2 activities not only enhances the confidence in the 2 assay procedures but also paves the way for design and development of potential chemopreventive agents. No apparent differences were obsd. in the **binding** affinities of the retinoids to **binding** proteins of a normal tissue or a tumor tissue. No correlation existed between the **binding** affinities of these retinoids to serum albumin and their biol. activity. Structure-activity relationships of the retinoids in relation to their **binding** affinities and biol. activities have been discussed.

IT 75664-66-3

RL: BIOL (Biological study)
(**binding** of, by retinoic acid-**binding** protein and serum albumin, neoplasm **inhibition** in relation to)

L53 ANSWER 26 OF 26 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1983:12895 HCAPLUS

DOCUMENT NUMBER: 98:12895

TITLE: Laboratory evaluation of flumethrin, a new synthetic pyrethroid for the control of one- and multi-host ticks

AUTHOR(S): Stendel, W.; Fuchs, R.

CORPORATE SOURCE: Inst. Chemother., Bayer A.-G., Wuppertal, D-5600, Fed. Rep. Ger.


SOURCE: VMR, Vet. Med. Rev. (1982), (2), 115-29

CODEN: VVMRDI; ISSN: 0341-9851

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



AB In-vitro tests and **controlled** animal trials with flumethrin (I, Bayticol) [69770-45-2] are reported. I **inhibits** egg deposition of normal sensitive and resistant one-host ticks of multi-host ticks, at <10 ppm. In animal tests, I at 30 ppm was 100% effective against Boophilus microplus, B. decoloratus and B. annulatus, even against strains resistant to all known tickicides and against multi-host Rhipicephalus appendiculatus, R. evertsi, Amblyomma hebraeum, A. variegatum, A. cajennense and Hyalomma truncatum. The onset of action is obsd. within a few h after treatment and the effect is irreversible. I is stable in dip-wash and does not strip. Animals tolerated concns. manifold the recommended ones and no irritating effects on skin or mucous membranes were obsd.

IT 69770-45-2

RL: BIOL (Biological study)
(tick **control** by)

=>

=>

=> select hit rn 132 ;select hit rn 152;select hit rn 153 1-26
ENTER ANSWER NUMBER OR RANGE (1-):1-
E20 THROUGH E21 ASSIGNED

ENTER ANSWER NUMBER OR RANGE (1-):1-
E22 THROUGH E22 ASSIGNED

E23 THROUGH E38 ASSIGNED

=> fil reg

FILE 'REGISTRY' ENTERED AT 10:33:52 ON 21 MAR 2002
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2002 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 19 MAR 2002 HIGHEST RN 401892-67-9
DICTIONARY FILE UPDATES: 19 MAR 2002 HIGHEST RN 401892-67-9.

TSCA INFORMATION NOW CURRENT THROUGH July 7, 2001

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Calculated physical property data is now available. See HELP PROPERTIES
for more information. See STNote 27, Searching Properties in the CAS
Registry File, for complete details:
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

The P indicator for Preparations was not generated for all of the
CAS Registry Numbers that were added to the H/Z/CA/CAPLUS files between
12/27/01 and 1/23/02. Use of the P indicator in online and SDI searches
during this period, either directly appended to a CAS Registry Number
or by qualifying an L-number with /P, may have yielded incomplete results.
As of 1/23/02, the situation has been resolved. Also, note that searches
conducted using the PREP role indicator were not affected.

Customers running searches and/or SDIs in the H/Z/CA/CAPLUS files
incorporating CAS Registry Numbers with the P indicator between 12/27/01
and 1/23/02, are encouraged to re-run these strategies. Contact the
CAS Help Desk at 1-800-848-6533 in North America or 1-614-447-3698,
worldwide, or send an e-mail to help@cas.org for further assistance or to
receive a credit for any duplicate searches.

=>

=>

=> d his 154

(FILE 'HCAPLUS' ENTERED AT 10:23:08 ON 21 MAR 2002)
SELECT HIT RN L32 1
SELECT HIT RN L52 1-
SELECT HIT RN L53 1-26

SELECT HIT RN L32 1-
 SELECT HIT RN L52 1-
 SELECT HIT RN L53 1-26

FILE 'REGISTRY' ENTERED AT 10:33:52 ON 21 MAR 2002

L54 19 S E20-E38

=>

=>

=> d ide can 154 1-19

L54 ANSWER 1 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 245742-21-6 REGISTRY

CN Pentanoic acid, 5-[2-[5-hydroxy-5-[1-(3-phenyl-2-propynyl)cyclobutyl]-1,3-pentadienyl]cyclohexylidene]- (9CI) (CA INDEX NAME)

OTHER NAMES:


CN ZK 158252

FS 3D CONCORD

MF C29 H36 O3

SR CA

LC STN Files: CA, CAPLUS



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

3 REFERENCES IN FILE CA (1967 TO DATE)

3 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 135:737

REFERENCE 2: 134:217509

REFERENCE 3: 131:267422

L54 ANSWER 2 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 173792-73-9 REGISTRY

CN 2,4-Pentadienoic acid, 3-methyl-5-[2-[(1E)-2-(2,6,6-trimethyl-1-cyclohexen-1-yl)ethenyl]-1-cyclohexen-1-yl]-, (2E,4E)- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 2,4-Pentadienoic acid, 3-methyl-5-[2-[2-(2,6,6-trimethyl-1-cyclohexen-1-yl)ethenyl]-1-cyclohexen-1-yl]-, (E,E,E)-

OTHER NAMES:

CN Ro 25-6603


FS STEREOSEARCH

MF C23 H32 O2

SR CA

LC STN Files: BIOSIS, CA, CAPLUS, TOXCENTER, USPATFULL

Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

5 REFERENCES IN FILE CA (1967 TO DATE)
5 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 126:305659

REFERENCE 2: 125:222217

REFERENCE 3: 124:278204

REFERENCE 4: 124:221563

REFERENCE 5: 124:166072

L54 ANSWER 3 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 153732-27-5 REGISTRY

CN Benzoic acid, 3-[2-[4-[2-[[4-chlorophenyl)sulfonyl]amino]ethyl]phenyl]-2-(3-pyridinyl)ethenyl]-, (Z)- (9CI) (CA INDEX NAME)


FS STEREOSEARCH

MF C28 H23 Cl N2 O4 S

SR CA

LC STN Files: CA, CAPLUS

Double bond geometry as shown.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1967 TO DATE)
1 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 120:217212

L54 ANSWER 4 OF 19 REGISTRY COPYRIGHT 2002 ACS
RN 153732-10-6 REGISTRY
CN Benzoic acid, 3-[2-[4-[2-[[4-chlorophenyl)sulfonyl]amino]ethyl]phenyl]-2-(3-pyridinyl)ethenyl]-, (E)- (9CI) (CA INDEX NAME)
FS STEREOSEARCH
MF C28 H23 Cl N2 O4 S
SR CA
LC STN Files: CA, CAPLUS

Double bond geometry as shown.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1967 TO DATE)
1 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 120:217212

L54 ANSWER 5 OF 19 REGISTRY COPYRIGHT 2002 ACS
RN 138254-19-0 REGISTRY
CN Benzoic acid, 4-[(1E)-2-[4-[(3-methylbutyl)thio]phenyl]-1-propenyl]- (9CI)
(CA INDEX NAME)
OTHER CA INDEX NAMES:
CN Benzoic acid, 4-[2-[4-[(3-methylbutyl)thio]phenyl]-1-propenyl]-, (E)-
OTHER NAMES:
CN MM 3986
CN SR 3986
FS STEREOSEARCH
MF C21 H24 O2 S
SR CA
LC STN Files: CA, CAPLUS, TOXCENTER

Double bond geometry as shown.



****PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT****

5 REFERENCES IN FILE CA (1967 TO DATE)
5 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 135:55572

REFERENCE 2: 125:316203

REFERENCE 3: 124:209

REFERENCE 4: 121:49616

REFERENCE 5: 116:34012

L54 ANSWER 6 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 123048-11-3 REGISTRY

CN 5-Heptenoic acid, 7-[(1S,2R,3S,4R)-3-[(1E,3R,4S)-3-hydroxy-4-phenyl-1-pentenyl]-7-oxabicyclo[2.2.1]hept-2-yl]-, (5Z)- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 5-Heptenoic acid, 7-[3-(3-hydroxy-4-phenyl-1-pentenyl)-7-oxabicyclo[2.2.1]hept-2-yl]-, [1S-[1.alpha.,2.alpha.(Z),3.alpha.(1E,3S*,4R*),4.alpha.]]-

CN 7-Oxabicyclo[2.2.1]heptane, 5-heptenoic acid deriv.

OTHER NAMES:

CN SQ 28568

FS STEREOSEARCH


MF C24 H32 O4

SR CA

LC STN Files: BEILSTEIN*, CA, CAPLUS, TOXCENTER, USPATFULL
(*File contains numerically searchable property data)

Absolute stereochemistry.

Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

6 REFERENCES IN FILE CA (1967 TO DATE)
 1 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA
 6 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 115:106005

REFERENCE 2: 115:22233

REFERENCE 3: 113:165424

REFERENCE 4: 112:235023

REFERENCE 5: 111:180726

REFERENCE 6: 111:167404

L54 ANSWER 7 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 119567-94-1 REGISTRY

CN Benzoic acid, 4-[(1E)-3-[4-(1,1-dimethylethyl)phenyl]-3-oxo-1-propenyl]-
 (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Benzoic acid, 4-[3-[4-(1,1-dimethylethyl)phenyl]-3-oxo-1-propenyl]-, (E)-

OTHER NAMES:

CN Ch 40

FS STEREOSEARCH


MF C20 H20 O3

SR CA

LC STN Files: BEILSTEIN*, CA, CAPLUS, TOXCENTER

(*File contains numerically searchable property data)

Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

7 REFERENCES IN FILE CA (1967 TO DATE)
7 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 125:316403
REFERENCE 2: 121:221053
REFERENCE 3: 120:95017
REFERENCE 4: 119:62406
REFERENCE 5: 115:111425
REFERENCE 6: 112:198609
REFERENCE 7: 110:165533

L54 ANSWER 8 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 119567-93-0 REGISTRY

CN Benzoic acid, 4-[(1E)-3-(3,4-diethylphenyl)-3-oxo-1-propenyl]- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Benzoic acid, 4-[3-(3,4-diethylphenyl)-3-oxo-1-propenyl]-, (E)-

OTHER NAMES:

CN Ch 20


FS STEREOSEARCH

MF C20 H20 O3

SR CA

LC STN Files: CA, CAPLUS, DDFU, DRUGU, TOXCENTER

Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

4 REFERENCES IN FILE CA (1967 TO DATE)
4 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 125:316403
REFERENCE 2: 116:252911
REFERENCE 3: 115:111425
REFERENCE 4: 110:165533

L54 ANSWER 9 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 110368-34-8 REGISTRY

CN Benzoic acid, 4-[(1E)-3-[3,4-bis(1-methylethyl)phenyl]-3-oxo-1-propenyl]-
(9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Benzoic acid, 4-[3-[3,4-bis(1-methylethyl)phenyl]-3-oxo-1-propenyl]-, (E)-

OTHER NAMES:

CN Ch 30

CN Ch 30 (retinoid)

FS STEREOSEARCH

DR 123008-50-4


MF C22 H24 O3

SR CA

LC STN Files: BEILSTEIN*, CA, CAPLUS, TOXCENTER

(*File contains numerically searchable property data)

Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

4 REFERENCES IN FILE CA (1967 TO DATE)

4 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 125:316403

REFERENCE 2: 115:111425

REFERENCE 3: 110:165533

REFERENCE 4: 107:126601

L54 ANSWER 10 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 110368-33-7 REGISTRY

CN Benzoic acid, 4-[(1E)-3-[3,5-bis(1,1-dimethylethyl)phenyl]-3-oxo-1-propenyl]- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Benzoic acid, 4-[3-[3,5-bis(1,1-dimethylethyl)phenyl]-3-oxo-1-propenyl]-, (E)-

OTHER NAMES:

CN Ch 55

FS STEREOSEARCH


MF C24 H28 O3

SR CA

LC STN Files: BEILSTEIN*, BIOSIS, BIOTECHNO, CA, CAPLUS, EMBASE, TOXCENTER, USPATFULL

(*File contains numerically searchable property data)

Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

49 REFERENCES IN FILE CA (1967 TO DATE)
 1 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA
 49 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 134:80645
 REFERENCE 2: 133:275967
 REFERENCE 3: 133:202727
 REFERENCE 4: 133:99541
 REFERENCE 5: 132:233218
 REFERENCE 6: 130:320477
 REFERENCE 7: 129:117505
 REFERENCE 8: 128:43515
 REFERENCE 9: 127:229396
 REFERENCE 10: 127:188955

L54 ANSWER 11 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 103455-29-4 REGISTRY

CN Benzeneacetic acid, .alpha.-(methoxymethylene)-2-[(1E)-2-phenylethenyl]-, methyl ester, (.alpha.E)- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Benzeneacetic acid, .alpha.-(methoxymethylene)-2-(2-phenylethenyl)-, methyl ester, (E,E)-


FS STEREOSEARCH

MF C19 H18 O3

SR CA

LC STN Files: CA, CAPLUS, MEDLINE, SPECINFO, TOXCENTER, USPATFULL

Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

14 REFERENCES IN FILE CA (1967 TO DATE)
 1 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA
 14 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 135:164018
 REFERENCE 2: 133:27880
 REFERENCE 3: 132:89965
 REFERENCE 4: 132:60924
 REFERENCE 5: 129:185980
 REFERENCE 6: 127:328208
 REFERENCE 7: 126:274931
 REFERENCE 8: 126:114577
 REFERENCE 9: 120:294381
 REFERENCE 10: 119:67877

L54 ANSWER 12 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 93060-40-3 REGISTRY

CN 5-Heptenoic acid, 7-[(1S,2R,3S,4R)-3-[(1E,3S)-3-cyclohexyl-3-hydroxy-1-propenyl]-7-oxabicyclo[2.2.1]hept-2-yl]-, (5Z)- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 5-Heptenoic acid, 7-[3-(3-cyclohexyl-3-hydroxy-1-propenyl)-7-oxabicyclo[2.2.1]hept-2-yl]-, [1S-[1.alpha.,2.alpha.(Z),3.alpha.(1E,3R*),4.alpha.]]-

CN 7-Oxabicyclo[2.2.1]heptane, 5-heptenoic acid deriv.

OTHER NAMES:

CN SQ 27427


FS STEREOSEARCH

MF C22 H34 O4

LC STN Files: AGRICOLA, BEILSTEIN*, BIOTECHNO, CA, CAPLUS, DDFU, DRUGU, EMBASE, PHAR, TOXCENTER
 (*File contains numerically searchable property data)

Absolute stereochemistry.

Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

4 REFERENCES IN FILE CA (1967 TO DATE)
4 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 121:73064

REFERENCE 2: 112:235023

REFERENCE 3: 104:230538

REFERENCE 4: 102:1050

L54 ANSWER 13 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 92843-52-2 REGISTRY

CN 5-Heptenoic acid, 7-[3-(3-hydroxy-3-phenyl-1-propenyl)-7-oxabicyclo[2.2.1]hept-2-yl]-, [1.alpha.,2.beta.(Z),3.beta.(1E,3R*),4.alpha.]-(9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 5-Heptenoic acid, 7-[3-(3-hydroxy-3-phenyl-1-propenyl)-7-oxabicyclo[2.2.1]hept-2-yl]-, [1.alpha.,2.beta.(Z),3.beta.(1E,3R*),4.alpha.]-(+.+.)-

CN 7-Oxabicyclo[2.2.1]heptane, 5-heptenoic acid deriv.


FS STEREOSEARCH

MF C22 H28 O4

LC STN Files: CA, CAPLUS

Relative stereochemistry.

Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1967 TO DATE)
1 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 101:204872

L54 ANSWER 14 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 87983-63-9 REGISTRY

CN 5-Heptenoic acid, 7-[(1R,2S,3R,4S)-3-[(1E,3S,4R)-3-hydroxy-4-phenyl-1-pentenyl]-7-oxabicyclo[2.2.1]hept-2-yl]-, (5Z)-rel- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

5-Heptenoic acid, 7-[3-(3-hydroxy-4-phenyl-1-pentenyl)-7-oxabicyclo[2.2.1]hept-2-yl]-, [1.alpha.,2.alpha.(Z),3.alpha.(1E,3S*,4R*),4.alpha.] -

CN 7-Oxabicyclo[2.2.1]heptane, 5-heptenoic acid deriv.

OTHER NAMES:

CN SO 28668

FS STEREOSEARCH


MF C24 H32 O4

LC STN Files: ANABSTR, BEILSTEIN*, BIOSIS, BIOTECHNO, CA, CAPLUS, DDFU,
DRUGU, EMBASE, IPA, MEDLINE, PHAR, PROMT, TOXCENTER, USPATFULL
(*File contains numerically searchable property data)

Relative stereochemistry.

Double bond geometry as shown.

Currently available stereo shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

14 REFERENCES IN FILE CA (1967 TO DATE)
14 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 136:694

REFERENCE 2: 135:205553

REFERENCE 3: 127:130997

REFERENCE 4: 121:73064

REFERENCE 5: 115:106005

REFERENCE . 6: 111:33370

REFERENCE 7: 109:184166

REFERENCE 8: 108:48997

REFERENCE 9: 108:30566

REFERENCE 10: 107:624

L54 ANSWER 15 OF 19. REGISTRY COPYRIGHT 2002 ACS

RN 75664-66-3 REGISTRY

CN Benzoic acid, 4-[(1E,3E)-2-methyl-4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3-butadienyl]- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Benzoic acid, 4-[2-methyl-4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3-butadienyl]-, (E,E)-

OTHER NAMES:

CN BASF 37400

CN SR 3983


CN SRI 2965-38

FS STEREOSEARCH

MF C21 H26 O2

LC STN Files: BEILSTEIN*, CA, CAPLUS, SPECINFO, TOXCENTER, USPATFULL
(*File contains numerically searchable property data)

Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

19 REFERENCES IN FILE CA (1967 TO DATE)

2 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

19 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 127:355043

REFERENCE 2: 126:311735

REFERENCE 3: 124:209

REFERENCE 4: 120:315175

REFERENCE 5: 120:182467

REFERENCE 6: 120:95771

REFERENCE 7: 120:95680

REFERENCE 8: 119:151651

REFERENCE 9: 119:62406

REFERENCE 10: 116:158918

L54 ANSWER 16 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 69770-45-2 REGISTRY

CN Cyclopropanecarboxylic acid, 3-[2-chloro-2-(4-chlorophenyl)ethenyl]-2,2-dimethyl-, cyano(4-fluoro-3-phenoxyphenyl)methyl ester (9CI) (CA INDEX NAME)

OTHER NAMES:

CN Bayticol

CN Bayticol Pour-on

CN Bayvarol

CN FCR 2769

CN Flumethrin

FS 3D CONCORD

MF C28 H22 Cl2 F N O3


CI COM

LC STN Files: AGRICOLA, ANABSTR, BEILSTEIN*, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CABA, CAPLUS, CBNB, CHEMCATS, CHEMLIST, CIN, CSCHEM, CSNB, DDFU, DRUGU, EMBASE, MEDLINE, MRCK*, NIOSHTIC, PROMT, TOXCENTER, USPATFULL; VETU

(*File contains numerically searchable property data)

Other Sources: EINECS**

(**Enter CHEMLIST File for up-to-date regulatory information)



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

115 REFERENCES IN FILE CA (1967 TO DATE)

5 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

116 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 136:133816

REFERENCE 2: 136:4861

REFERENCE 3: 135:340456

REFERENCE 4: 135:268702

REFERENCE 5: 135:222865

REFERENCE 6: 135:15451

REFERENCE 7: 134:371766

REFERENCE 8: 134:349373

REFERENCE 9: 134:174233

REFERENCE 10: 134:143283

L54 ANSWER 17 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 65199-97-5 REGISTRY

CN Benzoic acid, 4-[2-[4-(dimethylamino)phenyl]ethenyl]-3-nitro- (9CI) (CA INDEX NAME)


FS 3D CONCORD

MF C17 H16 N2 O4

CI COM

LC STN Files: BEILSTEIN*, CA, CAPLUS

(*File contains numerically searchable property data)



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

7 REFERENCES IN FILE CA (1967 TO DATE)

1 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

7 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 131:229682

REFERENCE 2: 112:37143

REFERENCE 3: 111:135189

REFERENCE 4: 110:24585

REFERENCE 5: 91:56115

REFERENCE 6: 89:42075

REFERENCE 7: 88:21879

L54 ANSWER 18 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 61059-47-0 REGISTRY


CN Benzenepropanoic acid, 4-(3-oxo-3-phenyl-1-propenyl)- (9CI) (CA INDEX NAME)

FS 3D CONCORD

MF C18 H16 O3

LC STN Files: BEILSTEIN*, CA, CAPLUS, IFICDB, IFIPAT, IFIUDB, USPATFULL

(*File contains numerically searchable property data)



****PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT****

4 REFERENCES IN FILE CA (1967 TO DATE)
4 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 87:167872

REFERENCE 2: 87:152001

REFERENCE 3: 87:135018

REFERENCE 4: 85:177244

L54 ANSWER 19 OF 19 REGISTRY COPYRIGHT 2002 ACS

RN 20118-38-1 REGISTRY

CN Benzoic acid, 4-(3-oxo-3-phenyl-1-propenyl)- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Benzoic acid, p-(2-benzoylvinyl)- (8CI)

OTHER NAMES:


CN 4-(2-Benzoylvinyl)benzoic acid

FS 3D CONCORD

MF C16 H12 O3

CI COM

LC STN Files: BEILSTEIN*, CA, CAPLUS, CASREACT, IFICDB, IFIPAT, IFIUDB,
SPECINFO, TOXCENTER, USPATFULL
(*File contains numerically searchable property data)



****PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT****

24 REFERENCES IN FILE CA (1967 TO DATE)
3 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA
24 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 135:5455

REFERENCE 2: 130:52857

REFERENCE 3: 129:261547

REFERENCE 4: 129:203719

REFERENCE 5: 128:322085

REFERENCE 6: 128:302106

REFERENCE 7: 127:347733

REFERENCE 8: 126:187485

REFERENCE 9: 126:105565

REFERENCE 10: 125:198514